Mental health medication story

A guide to talking about medication for mental health















Yarning about mental health
Trisia Nagal & Carolyn Thompson AlMbi N

Tricia Nagel & Carolyn Thompson, AlMhi NT











Inside your head is your brain.

Each person's brain is very different.

The brain is like our control centre.


The brain sends messages around the body.

Messages are carried by natural chemicals in the body — like water in a healthy river system.



In a healthy person, all the chemicals in the nervous system are balanced.

Like water in a healthy river system.







Some things can make the balance in our nervous system go all wrong.

Things like:

- using grog, gunja or other drugs
- sniffing petrol
- too much worries or stress.



Drugs, alcohol and stress can make the balance wrong. The person's balance is all wrong like a river system with dams or bad chemicals.







When the balance goes wrong in the nervous system, confused messages are sent around the brain and body.

When the balance is all wrong in the person's brain and body, they may get a mental illness.





Treatment for mental illness is Two Way —

Aboriginal way and Western medicine way working together









We can get back in balance by making changes.

Sometimes we need Western medicine.

Sometimes we need Aboriginal way. Sometimes both.





Social changes are about doing different things.

Like playing footy or going fishing.











Other social changes for better mental health might be to talk to Elders or to council or to clinic mob — or to mental health mob.









Psychosis







Depression







Anxiety





There are three main sorts of tablets for mental illness.

antipsychotics



antidepressants antianxiety tablets





Mental health medicine taken for **one** day will help balance for **one** day. Medicine taken **every** day will help balance **every** day.







Medicine can be hard to take regularly.

Put medication where you can see it so you remember to take it.





Take medication the same time each day — like when you get out of bed — so you remember to take it.









Use a dosette or Webster pack each day so you remember to take it.







There are three common early warning signs of getting sick with mental illness again:

- 1. Change of appetite
- 2. Worry or restlessness
- 3. Change of sleep.





If you notice early warning signs of getting sick again – make changes and get help quickly.





The three common trigger factors for getting sick again are:

- 1. Stopping treatment
- 2. Life troubles
- 3. Grog, gunja or other drugs.








If you have trigger factors or early warning signs of getting sick again – make changes and get help quickly.



Mental health medicine needs to be checked regularly by your doctor. Sometimes you need **more** for balance, sometimes you need **less**, sometimes you need to **change** medicine.

Use the dosette or Webster pack each day so you remember to take it. If you have trouble remembering tablets, mental health injections (needles) might help. Mental health injections can help balance for weeks.





Feedback invited – AIMhi NT Tricia.Nagel@menzies.edu.au Carolyn.Griffin@menzies.edu.au www.menzies.edu.au/AIMhi

Graphic Design: Inprint Design (08) 8201 3223 • www.inprint.com.au

Artwork: Tricia Nagel with valued assistance from Carolyn Thompson and Robert Mills

Photography: Mary Moloney, Robert Mills, Glen Norris, Tricia Nagel, Valerie Thompson, Neil Spencer, Sheree Cairney and Joseph Fitz

Copyright © 2008 Menzies School of Health Research (Menzies) ABN 70 413 542 847 Building 58, Royal Darwin Hospital Campus Rocklands Drive, Casuarina NT 0810, Australia.

All rights reserved. Except as provided in the Copyright Act 1968, this work may not be reproduced in any form without the written permission of Menzies School of Health Research

ISBN ISBN 978-1-921889-09-7

AlMhi NT is supported by DHCS and CRCATSIH

